## Remotet DS Vet Bolus

## Oxytetracycline Hydrochloride BP

Composition: Each Bolus contains Oxytetracycline Hydrochloride BP eq. to Oxytetracycline 1.0 gm.

**Description:** Oxytetracycline is a broad spectrum antibiotic highly active against a wide range of gram positive and gram negative organisms including most of the pathogenic bacteria of animals.

## Indication:

Cattle, Buffalo, Goat, Sheep and Horse: Remotet DS (Oxytetracycline) Bolus is indicated for anthrax, haemorrhagic septicaemia, black quarter, brucellosis, pyelonephritis, calf scour, infectious enteritis, navel ill, dysentery, pneumonia, pleuropneumonia, abscess, liver abscess, dermatophilosis, mastitis, malignant edema, peritonitis, leptospirosis, infectious calf diphtheria, infectious bovine keratoconjunctivitis, salmonellosis, actinomycosis, anaplasmosis, foot-rot, dehorning, castration etc. Remotet DS

Bolus is also indicated for secondary infections caused by virus such as foot and mouth disease (FMD), ephemeral fever etc.

**Dog and Cat:** Pneumonia, pleurisy, bronchitis, infectious enteritis and secondary infections in distemper. **Poultry:** Remotet DS Bolus is used for the prevention and treatment of different infections susceptible to Oxytetracycline, such as colibacillosis, salmonellosis, fowl cholera (pasteurellosis), infectious coryza, chronic respiratory disease (CRD) and infectious sinuvitis.

## **Dosage & Administration:**

Cattle and Buffalo: 1-2 Bolus per 50-100 kg body weight. Goat and Sheep: ½ Bolus per 10-20 kg body weight.

Horse: ½ Bolus per 75 kg body weight.

Poultry: 1 Bolus per 4 liters of Drinking water or 1 Bolus per 10 kg feed.

Or, as prescribed by the registered veterinary doctor.

 $\textbf{Contra-Indication:} \ \ \text{Do} \ \ \text{not} \ \ \text{administer} \ \ \text{those} \ \ \text{animals} \ \ \text{which} \ \ \text{are} \ \ \text{hypersensitive} \ \ \text{to} \ \ \text{tetracycline} \ \ \text{or} \ \ \text{those}$ 

animals which have  $% \left( 1\right) =\left( 1\right) \left( 1\right)$ 

serious renal and liver impairments.

Withdrawal time:

Cattle: Meat: 15-22 days,

Poultry: Meat: 05 days.

Lactating Animal: Milk: 2 days

Storage: Protect from light, store in a cool & dry place and keep out of reach of children.

Presentation: Each box contains 3 X 4 Bolus in Alu-Alu Strips.