Regenta 10%® Injection

Gentamicin Sulphate BP

Composition:

Each ml contains Gentamicin Sulphate BP equivalent to Gentamicin 100 mg.

Indications: Regenta 10% Injection is effective against various types of bacterial diseases such as:

Poultry: Salmonellosis, Colibacillosis, Swollen Head Syndrome, Mycoplasmosis.

Goat & Sheep: Colibacillosis, Pasteurellosis, Mastitis, Pneumonia.

Cow, Buffalo: Colibacillosis, Mastitis, Pneumonia, Metritis.

Horses: Pleuropneumonia, Enteritis, Polyarthritis, Endometritis.

Dosage & Administration:

Regenta 10% Injection preferably to be given twice for 1st day and afterwards once daily for 3-5 days at following dose:

Livestock: 1 ml per 25 kg body weight

Poultry: 0.05 ml – 0.1ml per kg body weight

Or, as prescribed by the registered veterinary doctor

Side Effects: When Regenta 10% Injection is given in the recommended doses noadverse effects are to be expected but prolonged treatment with higher doses can cause possible renal damage, ototoxicity, neuromuscular paralysis or hypocalcaemia.

Contra-indication: Regenta 10% Injection is contraindicated for parenteral administration in renal insufficiency.

Withdrawal Period:

Meat: 15 days (For kidneys, liver and the site of injection: 45 days).

Milk: 03 days.

Storage: Protect from light, store in a cool & dry place and keep out of reach of children.

Presentation: 10 ml, 25 ml, 50 ml, 100 ml & 200 ml Vial